

NIR and NMR Soybean Composition Database

*I.C. Baianu, **Marvin Paulsen and ***Randall L. Nelson

* AFC-NMR and NIR Microspectroscopy Facility

** Agricultural Engineering Department

*** National Soybean Laboratory and Crop Sciences Department

College of ACES, University of Illinois, Urbana, Illinois 61801,
USA.

Abstract

Our NIR and NMR data acquired since 2002 with cutting edge instrumentation and novel NIR calibrations will allow the development of an *Internet Database of Soybean Composition* for 15,000 Soybean Accessions from the National Soybean Collection at the University of Illinois, containing the data obtained from over 80,000+ spectroscopic NIR and NMR measurements for: *proteins, amino acids, oils, fatty acids, isoflavones, carbohydrates, fiber, moisture* in intact soybean seeds both in bulk and single soybean seed samples.

This novel **Soybean Composition Database** will also include more than 12,000 NIR measurements on soybeans from the **International Soybean Germplasm Collection**, such as those received from Peking at the National Soybean Collection.

XLS files of our novel spectroscopic data are currently available for *all 80,000+* NIR and FT-NMR measurements—such data will be made available from an ultra-fast and secure supercomputer server utilizing the current version of the *Scientific-Linux* OS-based software.

1. FT-IR and FT-NIR Microspectrometers

A microspectrometer is defined as a combination of a spectrometer and a microscope that has both spectroscopic and imaging capabilities. Such an instrument is capable, for example, of obtaining visible images of a sample using a CCD camera, and chemical images with an NIR detector. Chemical images are then employed for sophisticated quantitative analyses. The results reported in this chapter for soybean seeds and embryos were obtained with FT-IR and -NIR spectrometers made by the PerkinElmer Co. (Shelton, CT, USA). The FT-NIR (NTS model) spectrometer was equipped with an integrating sphere accessory for diffuse reflectance. The FT-IR or -NIR spectrometers were, respectively, attached to microscopes for the IR region (Spotlight 300) or NIR region (NIR Autoimage), as illustrated in Fig. 3.2.1 and Fig. 3.2.2. Each spectrometer has an internal desiccant compartment to remove the water vapor and the carbon dioxide from air that may interfere with the spectrum of a sample. Apart from the improved resolution and acquisition time, these instrument models, offer increased sensitivity and also allow the transfer of spectra to different instruments of similar design. The two microspectrometers are each equipped with two cassegrain imaging objectives and a third cassegrain before the NIR detector in order to improve focus and sensitivity, as shown in Fig. 3.2.3.

- ☐ Introduced in 2002 by the PerkinElmer Co. (Shelton, CT, USA) for high-resolution studies

- ☐ Employed for our mid-IR Microspectroscopy and Chemical Imaging investigations of thin sections



Fig. 1.1. FT-IR Microspectrometer (Spotlight 300) introduced by the PerkinElmer Co.

- ❑ Introduced in 2002 by PerkinElmer Co. (Shelton, CT, USA) for high-resolution studies
 - ❑ Employed for our NIR Microspectroscopy and Chemical Imaging investigations of soybean seeds and embryos
-




Fig. 1.2. FT-NIR Microspectrometer (AutoImage) made by the PerkinElmer Co.




Fig. 1. 3. A simplified diagram of the reflection mode of operation for the AutoImage

FT-NIR Microspectrometer, manufactured by the PerkinElmer Co.



Fig.1.4. The FCS Alba™ Microspectrometer System manufactured by ISS Co., (Urbana, Illinois, USA). The inverted, epi-fluorescence microscope shown in the figure is the Nikon TE-300 –special Model, that has available *both* a back illumination port and a left-hand side port. The PC employed for data acquisition, storage and processing is located behind the instrument, as is the laser illumination source (not visible in the figure).

2.1 High-Resolution NMR Method for Oil Determination

The technique applied to obtain the oil content in soybean embryos was simple one-pulse, High-Resolution (HR) NMR (11). The HR-NMR technique was explained in Section 3.4 of Chapter 1x. A Varian U-400 NMR instrument was employed for oil measurements; the selected 90 deg pulse width was 19.4 μ s and the ^1H NMR signal absorption intensity was recorded with a 4 s recycling interval to avoid saturation.

2.1.1 Oil Determination in Somatic Soybean Embryos by ^1H High-Resolution Nuclear Magnetic Resonance

High-resolution ^1H NMR measurements of oil were carried out with an U400 MHz NMR spectrometer as described in Section 3. A complete, nanoliter (nL) range oil calibration is reported here

for soybean somatic embryos with a measurement precision of $\sim \pm 0.1\%$ oil (calculated as a percentage of oil from the total, wet embryo weight). These are the first reported results of oil determination by high-resolution NMR in soybean somatic embryogenic cultures. From a 105-sample lot investigated, 89 were considered valid for further growth and selection. A summary of our is presented in Tables 4.3.1 to 4.3.4.

The corresponding quantity of oil in the embryos was calculated from a linear fitting of the HR-NMR data in our standard plot of soybean oil shown in Fig.4.3.1 with the following equation:

$$x = \frac{y + 0.0092}{0.0029},$$

where y was the normalized value of the oil peak integral from the experimental NMR spectrum of each sample (Fig.2.1.). The ratios of a chemical group proton signal other than water protons, with respect to the water proton signal--and the wet mass of the sample-- were then compared with the oil standard plot in order to estimate the quantity of oil present in soybean embryos.




Fig. 2.1. The soybean oil standard plot for 400 MHz ^1H NMR measurements on the

Varian U400. The probe was a Nalorac 5 mm QUAD for high-resolution

3. Soybean Correlation Data

The 99% linear correlation between the ^{13}C SS-NMR and the corresponding Dual DA-NIRS oil and protein measurements on the same samples of well-defined soybean accessions from the USDA Soybean Germplasm Collection at UIUC, suggests that both techniques are suitable for the non-destructive, practical determination of both oil and protein content of soybean flours.




Fig. 3.1. VACP ^{13}C SS-NMR Measurements of Protein Content in Soybean Flours and their direct, Linear Correlation with the Corresponding NIR Data.




Fig. 3.2. 1PDNA ^{13}C SSNMR of Oil Content of Soybean Flours and their direct linear correlation with the corresponding NIR data.




Fig.3.3. Protein-Oil Inverse Correlation of 5,000 Soybean Samples of Experimental Lines at UIUC.

4. EXCEL Data Tables of Soybean Accessions Composition

Table 4.1. VACP ¹³C SS-NMR Measurements of Protein Content of Soybean Flours and Their Direct Correlation with the NIR Data (as shown in Section 2.6).

Soybean Seed ID	M2123	W2101	W2403	M309	M285	96-960A2-2687	W1228	96-959A6-1447	LG00-13523	LG00-13251
% protein by NIR	37.0	39.1	40.8	42.7	43.8	46.2	49.6	52.6	55.0	56.7
% protein by NMR	37.1	40.0	41.5	42.0	44.0	47.0	49.0	53.1	55.0	58.0

Table 4.2. Range of constituents for 66 selected soybean samples selected as standards for Fatty Acid NIR Calibrations (data is courtesy of the USDA Peoria Laboratory). (Soybean lines are identified as: Stoneville 1999, MG V–VIII. USDA Germplasm, National Research Center, Urbana, IL.)

Simple Statistics	Dry Protein	Dry Oil	Moisture	%16:0	% 18:0	% 18:1	% 18:2	% 18:3
Mean	47.61	17.11	5.86	11.91	3.24	21.19	55.42	8.24
Stdev	2.03	1.21	0.17	0.70	0.49	2.51	2.08	1.03
Max	53.74	19.32	6.19	14.10	4.35	27.63	60.74	10.93
Min	43.80	13.70	5.48	10.54	2.33	14.24	51.09	6.32

Table 4.3. HR-NMR results for the Control (0 mM EMS) Soybean Embryo Group.

EMS solution conc. (mM)	Sample ID	Number of transients	Embryo mass (mg)	Normalized oil signal integral in the 3 ppm-0 ppm region	Quantity of oil in embryos (nL/mg)	Percentage of Oil in wet embryos (%)
0	6-29-0A2-10-5	500	9.80	0.0357	18	1.7
0	6-23-0A2-10-1	500	10.90	0.0269	14	1.3
0	7-11-0A2,0C2,8	500	10.79	0.0255	13	1.2
0	5-14-4HR-W0	500	7.82	0.0232	12	1.1
0	6-19-0A2-10-15	500	8.90	0.0196	11	1.0
0	4-3-0A6	500	20.52	0.0162	10	0.9
0	6-19--0B2-11-13	500	10.30	0.0153	9	0.8
0	6-19-0B1-10-15	500	11.20	0.0141	9	0.8
0	7-25-0A5,0B8,0C7	500	21.01	0.0119	8	0.7
0	6-26-0B5-12-4	500	14.80	0.0083	6	0.6
0	6-23-0B3-9-15	500	21.50	0.0072	6	0.6
0	6-26-0B1-12-4	500	24.80	0.0052	5	0.5
0	6-26-0B4-12-4	500	26.70	0.0040	5	0.5
0	6-23-0A3-11-2	500	19.50	0.0010	3	0.3

Table 4.4. HR-NMR results for the 1 mM EMS group.

EMS solution conc. (mM)	Sample ID	Number of transients	Embryo mass (mg)	Normalized oil signal integral in the 3 ppm-0 ppm region	Quantity of oil in embryos (nL/mg)	Percentage of Oil in wet embryos (%)
1	4-3-1A10,11,1C7	500	7.61	0.0598	27	2.5
1	6-23-1A2-10-15	500	6.80	0.0451	21	1.9
1	6-23-1B4-10-15	500	6.70	0.0290	15	1.4
1	7-11-1A8,1B7	500	14.79	0.0254	13	1.2
1	7-25- 1A10,1B2,6,7	500	10.91	0.0241	13	1.2
1	7-25-1C3,9	500	12.56	0.0214	12	1.1
1	6-19-1A5-10-5	500	11.40	0.0192	11	1.0
1	4-3-1A5,6,11- 1B10	500	18.58	0.0137	9	0.8
1	6-19-1A7-11-2	500	11.80	0.0127	8	0.7
1	4-3-1A5-9-17-s	500	57.82	0.0067	6	0.6
1	6-19-1A12-11-13	500	11.90	0.0065	6	0.6
1	6-19-1A6-11-2	500	12.60	0.0013	3	0.3

Table 4.5. Average values of oil content (wet %) in somatic embryogenic cultures of soybean samples, measured by 1PULSE ^1H NMR experiments, and variation range.

Embryo Oil Average Values				
0 mM EMS Oil, %	1 mM EMS Oil,%	3 mM EMS Oil,%	10 mM EMS Oil,%	30 mM EMS Oil,%
0.8	1.1	0.7	1.2	0.9
Ranges				
3-17	3-25	4-11	4-17	3-25

Table 4.6.

Mean values and variances of protein and oil for 17 Bulk Soybean groups.

Soybean groups and instruments used	Sample number	P-mean	P-var	O-mean	O-var
Germplasm98, DA7000	400	46.2	13.55	19.6	2.23
MAPIII97-mix, F3, DA7000	496	43.8	5.53	21.4	1.50
MAPIII98-mix, F3, DA7000	529	45.1	6.56	20.1	1.73
MAPIII97-Bell, F3, IM9100	707	47.3	3.84	19.7	1.25
MAPIII97-Fisher, F3, IM9100	704	46.7	3.85	20.0	1.03
MAPIII97-EG1000, F3, IM9100	703	43.0	3.80	21.9	0.95
OPMAP98, F3, IM9100	379	44.5	6.19	20.8	2.54
Germplasm99, ZX-50	961	44.9	2.61	19.4	0.96
99ProSel-F3, ZX-50	72	51.2	3.69	16.7	0.28
99ProSel-F4, ZX-50	206	49.1	4.40	17.7	0.90
OPMAP99-Hume, F3, ZX-800	495	47.2	6.83	17.9	2.76
OPMAP99-Bell, F3, ZX-800	490	47.0	6.91	17.7	3.07
99ProGen, F5, ZX-800	362	49.3	6.31	16.9	1.71
2000YLDMAP-Hume, F6, ZX-800	380	42.9	1.68	22.0	0.55
2000YLDMAP-MIV, F6, ZX-800	380	42.0	2.70	22.4	0.75
2000 ProSel and BCPro, F8, ZX-800	79	51.3	3.67	16.5	1.14
2000 ProSel, F9, ZX-800	24	54.6	0.99	14.2	0.27

Note 1: P-mean, P-var, O-mean, O-var represent protein mean, protein variance, oil mean, oil variance, respectively.

5. NIRS Basic Results

Figure 5.1. Pure Component FT-NIR Spectra of Major Soybean Constituents: Protein, Oil, Moisture




Figure 5.2.

Comparison of DA-NIRS Spectra of Single Soybean Seeds with Bulk Sample




Figure 5.3. NIR Predicted vs Reference Protein Values by Nicolet
Antaris Single Seed Calibration




Figure 5.4. NIR Predicted vs Reference Oil Values by Nicolet Antaris Single Seed Calibration




Table 5.1.

**Correlation Coefficients (R) and Standard Errors of
Cross Validation (SECV) for Single Seeds Analysis on
the DA-7000 Instrument**

Component	Number of Factors	R	SECV
Protein	15	98.5%	1.1
Oil	16	98.5%	0.5
Moisture	16	99.0%	0.3

Table 5.2.

Correlation Coefficients (R) and Standard Error of Cross Validation (SECV) for Single Seed Analysis on the Nicolet Antaris FT-NIR Instrument

Component	Number of Factors	R	SECV
Protein	14	99.2%	0.77
Oil	14	99.0%	0.42
Moisture	11	99.1%	0.27

Table 5.3.

Correlation Coefficients (R) and Standard Errors of Cross Validation (SECV) for Single Seeds Analysis on the DA-7000 Instrument

Component	Number of Factors	R	SECV
Protein	15	98.5%	1.1
Oil	16	98.5%	0.5
Moisture	16	99.0%	0.3

Figure 5.5.

Protein-Oil Inverse Correlation for 120 Single Seed Soybean Samples




Figure 5.6.




Figure 5.7.

**Protein-Oil Inverse Correlation for 4500
Map3 (F3 Generation, Harvested in '97 & '98)
Bulk Soybean Samples**




Figure 5.8.

Pure Component FT-NIR Spectra of Carbohydrate Soybean Constituents: Small sugars, and Fiber





Figure 5.9.

Protein Calibration for Bulk Soybean Analysis on the Spectrum One NTS FT-NIR Instrument



65 calibration standards, 20 grams for each standard, 8.9mm NIR beam size

Source: Soybean NIR Database, UIUC

Protein Calibration for Bulk Soybean Analysis on the DA-7000 Diode Array NIR Instrument




Moisture Calibration for Bulk Soybean Analysis on the DA-7000 Diode Array NIR Instrument



Figure 5.12.

Moisture Calibration for Bulk Soybean Analysis on the Spectrum One NTS FT-NIR Instrument



65 calibration standards, 20 grams for each standard, 9mm NIR beam size

Source: Soybean NIR Database, UIUC

Table 5.4.

**Calibration Results for Protein and Oil
Analysis with the Perten DA-7000, Dual
Diode-Array DA-NIR Instrument**

Components	Protein		Oil	
	Bulk Sample	Single Seeds	Bulk Sample	Single Seeds
SECV	0.1	1.1	0.1	0.5
R	99.9%	98.5%	99.9%	98.5%

Source: Soybean NIR Database, UIUC

Table 5.5.

**Bulk Soybean Calibration with 65 Standards
for Protein, Oil, and Moisture Analysis**

Developed with Data from the SpectrumOne NTS FT-NIR Spectrometer

Component	Number of Factors	R	SECV	SEP
Protein	13	99.9%	0.26	0.33
Oil	15	99.9%	0.13	0.23
Moisture	15	99.9%	0.17	0.30

R : **Correlation Coefficient**

SECV: **Standard Error of Cross Validation**

SEP: **Standard Error of Prediction**

Source: Soybean NIR Database, UIUC

Table 5.6.

Detrimental Effects of Light Scattering on the Accuracy of NIR Analysis for Whole Soybeans

(measured with the FT-NIR, SpectrumOne NTS Spectrometer)

Component	R		SECV	
	No MSC	MSC	No MSC	MSC
Protein	99.5	99.9	0.63	0.26
Oil	99.3	99.9	0.29	0.13
Moisture	99.8	99.9	0.26	0.17

R: Correlation coefficient

SECV: Standard Error of Cross Validation

(Tested with 65 bulk whole soybean standards)

Source: T. You, 2006

6. AMINO ACID COMPOSITION OF SOYBEAN SEEDS: NIR CALIBRATIONS

Table 6.1.

Correlations of Amino Acids with Crude Protein in Soybeans

A.A.	R ²	R
ASX	0.634	0.796
THR	0.005	0.071
SER	0.578	0.760
GLX	0.690	0.831
PRO	0.480	0.693
GLY	0.628	0.792
ALA	0.619	0.787
VAL	0.640	0.800
ILE	0.631	0.794

A.A.	R ²	R
LEU	0.644	0.802
TYR	0.536	0.732
PHE	0.474	0.689
HIS	0.499	0.706
LYS	0.652	0.807
ARG	0.687	0.829
MET	0.548	0.740
CYS	0.498	0.706
M+C	0.538	0.733

Table 6.2.

Amino Acids **Highly-Correlated** with the Dry Protein Content:

- Histidine: **R = 0.93**
- Arginine: **R = 0.90**
- Glx: **R = 0.88**
- Valine: **R = 0.87**
- Leucine: **R = 0.85** → *borderline*

=====

and the **imino acid** Proline: **R = 0.87**

Figure 6. 1.

Histidine: His vs. % Dry Protein




Figure 6.2.

Glutamine plus Glutamic Acid, "GLX", as
Total Dry Weight % vs. Dry Soybean Protein %




Table 6.3.

VIP-Data		Example:		
10DataPoints	TOTAL:	3,816data-points		
Prot	Moist	Oil	DP	DP calc
	DO calc			
35.57	11.10	20.50	40.01	40.01
	23.06			
36.91	10.00	20.76	41.00	41.01
	23.07			
41.75	10.41	17.06	46.60	46.60
	19.04			
41.44	10.30	10.02	46.20	46.20
	11.17			
41.10	10.16	18.60	45.75	45.75
	20.70			
36.71	9.12	20.60	40.40	40.39
	22.67			
38.03	10.84	20.24	42.65	42.65
	22.70			
39.52	10.44	19.34	44.13	44.13
	21.59			
40.17	9.04	18.80		44.16
	20.67			
40.67	10.25	18.30	45.31	45.31
	20.39			

Figure 6.3.

FT-NIR Spectra of Five Major Soybean Components

Collected on the Perkin-Elmer SpectrumOne NTS FT-NIR Spectrometer





Figure 6.4.


Predicted vs. Reference Concentrations of Component C1 for our PLS-1 Simulation Study



Source: T. You, 2006

Figure 6.5. Matrix NIR Analysis setup for three AA components.

Computer Simulation of PLS-1 Algorithm
with 3 Components (%)




Adapted from T. You, 2006

7. NEW SOYBEAN SEED NIR CALIBRATIONS

Figure 7.1.

A new and Improved Set of 124 Bulk Soybean Standard Samples:

Protein-Oil Inverse Correlation for the Year 2003 Calibration Standard



* 124 Standard samples were selected with a wide range of Protein and Oil concentrations that were uniformly distributed in 0.5% concentration steps for Protein, and in 0.2% steps for Oil. Source: Soybean NIR Database, UIUC

REFERENCES

- AACC. 1995. *Approved Methods of Analysis*, 9th Ed. The American Association of Cereal Chemists. St. Paul, Minnesota.
- AOAC International. 1995. *Official Methods of Analysis*, 16th ed., AOAC International, Gaithersburg, Maryland.
- AOCS. 1998. *Official Methods and Recommended Practices of the AOCS* – 5th Ed. The American Oil Chemists' Society. Champaign, Illinois.
- AOCS. 2002. Additions and Revisions to the Official Methods and Recommended Practices of the AOCS. AOCS Press. Champaign, IL. AK4-95.
- Augustine, M. and Baianu, I.C. 1984. High-Resolution ¹³C NMR and Ion Exchange Chromatography Amino Acid Profiling of Corn and Soybean Protein Hydrolysates. *FASEB Proceed.* 1125.
- Augustine, M.E., Baianu, I.C. 1986. High-Resolution ¹³C Nuclear Magnetic Resonance Studies of Maize Proteins. *J. Cereal Science.* 4: 371-378.
- Baianu, I.C. and H. Förster. 1980. Cross-Polarization, High-Field ¹³C NMR Techniques for Studying Physicochemical Properties of Wheat Grains and Wheat Proteins., *J. Applied Biochemistry*, 2: 347-355.
- Baianu, I.C. 1981. Carbon-13 and Proton NMR Studies of Wheat Proteins in Solution. *J. Sci. of Food & Agriculture*, 32: 309-313.
- Baianu, I.C., et al. 1990. Ch8. In "*NMR Applications in Biopolymers*". Edited by J.W. Finley et al., Plenum Press, New York.
- Baianu, I.C. 1992. Ed. "*Physical Chemistry of Food Processes. Vol. 1: Fundamental Aspects.*" Van Nostrand Reinhold. New York.
- Baianu, I.C., Pessen, H., Kumosinski, T.F. 1993. Ed. "*Physical Chemistry of Food Processes. Vol. 2: Advanced Techniques, Structures, and Applications.*" Van Nostrand Reinhold. New York.
- Baianu, I.C., You, T., Guo, J., Nelson, R. L. 2002. Calibration of Dual Diode Array and Fourier Transform Near Infrared Reflectance Spectrometers for Composition Analysis of Single Soybean Seeds in Genetic Selection, Cross-Breeding Experiments, Proceedings for the 9th Biennial Conference of the Cellular and Molecular Biology of the Soybean, August 11-14, 2002, 508.
- Baianu, I.C., Costescu, D., You, T. 2002. Novel Techniques for Microspectroscopy and Chemical Imaging Analysis of Soybean Seeds and Embryos, Proceedings for the 9th Biennial Conference of the Cellular and Molecular Biology of the Soybean, August 11-14, 2002, 509.
- Baianu, I.C., You T., Costescu, D.M., Lozano, P.R., Prisecaru, V. and Nelson, R.L. 2004a. Chapter 11 "High-Resolution Nuclear Magnetic Resonance and Near-Infrared Determination of Soybean Oil, Protein and Amino Acid Residues in Soybean Seeds", Luthria, D. Ed., *Oil Extraction and Analysis—Critical Issues and Comparative Studies*, AOCS Press. Champaign, IL. p193-240.
- Baianu, I.C., You T., Costescu, D.M., Lozano, P.R., Prisecaru, V. and Nelson, R.L. 2004b. Chapter 12 "Near-Infrared Microspectroscopy, Fluorescence Microspectroscopy, Infrared Chemical Imaging and High-Resolution Nuclear Magnetic Resonance Analysis of Soybean Seeds, Somatic Embryos and Single Cells, Luthria, D. Ed., *Oil Extraction and Analysis—Critical Issues and Comparative Studies*, AOCS Press. Champaign, IL. p241-273.
- Baianu, I.C., Korban, S.S., Costescu, D., You, T., Lozano, P., Hofmann, N.E. 2004c. Fourier Transform Near Infrared Microspectroscopy, Infrared Chemical Imaging, High-Resolution Nuclear Magnetic Resonance and Fluorescence Microspectroscopy Detection of Single Cancer Cells and Single Viral Particles. *Cern. Ext-2004-069*.
- K.A., and Lyman, C.M. 1948. Essential amino acid composition of soy bean meals prepared from twenty strains of soy beans. *J Biol. Chem.* 6: 28-36.

- Bardet, M., Foray, M.F. Bourguignon, J., and Krajewski, P. 2001. Investigation of Seeds with High-Resolution Solid-State ^{13}C NMR. *Magnetic Resonance in Chemistry*, 39: 733-738.
- Barton F.E. 2002. Theory and Principles of Near-Infrared Spectroscopy, *Spectroscopy Europe*, 14(1): 12-18.
- Belitz, H.D. and Grosch, W. 1987. *Food Chemistry*, Springer-Verlag, Berlin.
- Ben-Gera, I., Norris, K.H. 1968. Determination of Moisture Content in Soybeans by Direct Spectrophotometry. *Isr. J. Agric. Res.* 18: 125-132.
- Bardet, M., Foray, M.F., Bourguignon, J., and Krajewski, P. 2001. Investigation of Seeds with High-Resolution Solid-State ^{13}C NMR. *Magnetic Resonance in Chemistry*, 39: 733-738.
- Birth, G. S. 1986. The Light Scattering Characteristics of Ground Grains. *Intern. Agrophys.* 2(1): 59-67.
- Brim, C.A. and Burton, J.W. 1979. Recurrent Selection in Soybeans (*Glycine max*): 2. Selection for Increased Percent Protein in Seeds. *Crop Sci.* 19(4): 494-498.
- Burton, J.W., Wilson, R.F., Brim, C.A., and Rinne, R.W. 1989. Registration of Soybean Germplasm Line with Modified Fatty Acid Composition of Seed oil. *Crop Sci.* 29: 1583.
- Burton, J.W., and Wilson, R.F. 1994. Registration of N88-480, a Soybean Germplasm Line with High Concentration of Oil in Seeds. *Crop Sci.* 34: 313-314.
- Buning, P.H. and Diller, M. 2000. Rapid Analysis of Foods Using Near-Infrared Spectrometry (NIRS): Development, Use and New Possibilities. *Ernahrungs Umschau.* 47(1): 15-20.
- Buslov, D.K., Nikonenko, N.A., 1997. Regularized Method of Spectral Curve Deconvolution. *Applied Spectroscopy*, 51(5): 666-672.
- Buslov, D.K., Nikonenko, N.A., Sushko, N.I. and Zhbankov, R.G. 1999A. Analysis of the Results of α -D-glucose Fourier Transform Infrared Spectrum Deconvolution: Comparison with Experimental and Theoretical Data. *Spectrochimica Acta Part A.* 55: 229-238.
- Buslov, D.K., Nikonenko, N.A., Sushko, N.I. and Zhbankov, R.G. 1999B. Deconvolution of Fourier Transform Infrared Spectrum of α -D-Galactose: Comparison with Experimental and Theoretical Data. *Spectrochimica Acta Part A.* 55: 1101-1108.
- Cameron, A.G. 1967. An Assessment of the Potential Application of the Method of Attenuated Total Reflectance (ATR) Infrared Qualitative Analysis to Food Materials. *J. Food Technology.* 2:223.
- Carrao, P.M., de-Goes, F.S. and Kikuchi, A. 2002. Extraction Time for Soybean Isoflavone Determination. *Brazilian Archives Bio. Technol.* 45 (4): 515-518.
- Carr, H.Y., and Purcell, E.M. 1954. Effects on Diffusion on Free Precession in Nuclear Magnetic Resonance Experiments. *Physical Review*, 94 (3): 630-638.
- Cinier, R., Cuilment, J. 1998. PLS Calibration for Precise and Practical Quantitative Analysis of Simple Solutions Using NIR Spectroscopy. *Bruker Report*, 146:6-9.
- Coblentz, W.K. 1982. A Study in the Near Infrared Reflectance Spectra of Proteins and Related Compounds. MS thesis. Pennsylvania State University.
- Costescu, D.M. 2003. MS Thesis, University of Illinois at Urbana-Champaign, Urbana, Illinois.

- Delwiche, S.R., 1995. Single Wheat Kernel Analysis by Near-Infrared Transmittance: Protein Content. *Cereal Chem.* 72: 11-16.
- Derome, A.E. 1987. *Modern NMR Techniques for Chemistry Research*, p. 129, Pergamon Press, New York.
- Dev, S.B., Rha, C.K. and Walder, F. 1984. Secondary Structural Changes in Globular Protein Induced by a Surfactant: Fourier Self-Deconvolution of FT-IR Spectra. *J. Biomolecular Structure & Dynamics.* 2(2): 431-442.
- Diaspro, A., and Robello, M. 1999. Multi-Photon Excitation Microscopy to Study Biosystems. *European Microscopy and Analysis*, 5: 5-7.
- Diem, M. 1993. Ed. *Introduction to Modern Vibrational Spectroscopy*. John Wiley & Sons. New York, NY.
- Doty, D. M., Bergdoll, M. S., Nash, I. A., and Brunson, A. M. 1946. *Cereal Chem.* 23:199.
- Eigen, M. and Rigler, R. 1994. *Proc Natl. Acad. Sci. USA.* **91**: 5740.
- Fahrenfort, J. 1961. Attenuated Total Reflection. A New Principle for the Production of Useful Infrared Reflection Spectra of Organic Compounds. *Spectrochim. Acta* 17: 698
- Feynman, R.P. 1963. "*Lectures on Physics*", Vol.3, Addison-Wesley Publishing Company, Reading, MA.
- Fraenkel-Conrat, H, Cooper, M., 1944. The Use of Dyes for the Determination of Acid and Basic Groups in Proteins, *J. Biol. Chem.*, 154: 239-246.
- Galactic Industries Corporation. 1996. GRAMS/32 User's Guide: PLS plus/IQ for GRAMS/32 and GRAMS/386, Chapters 1-4.
- Geladi, P., MacDougall, D., Martens, H. 1985. Linearization and Scatter-Correction for Near- Infrared Reflectance Spectra of Meat. *Applied Spectroscopy.* 39(3): 491-500.
- Goldflus, F., Ceccantini, M., and Santos, W. Amino Acid Content of Soybean Samples Collected In Different Brazilian States: Harvest 2003/2004. *Rev. Bras. Cienc. Avic.* [Online]. 2006, 8(2): 105-111.
- Guo, J., You T., Baianu, I.C., Nelson, R.L. 2002. Evaluation of Four Dispersive NIR Instruments and Methodology Development for Soybean Composition Analysis in Genetic Selection and Breeding Programs, Proceedings of the China and International Soy Conference and Exhibition 2002 (CISCE 2002), November 6-9, 2002, 412-413.
- Guo, J. 2004. Development of Dispersive and Fourier Transform Near Infrared Spectroscopy Methodology for Food and Edible Seed Analysis. PhD Thesis, University of Illinois at Urbana-Champaign, Urbana, Illinois.

- Kuiken, Klempir J. M. 1999. Nuclear Magnetic Resonance Characterization of Soybeans for Oil Content Measurement. MS Thesis, University of Illinois at Urbana-Champaign, Urbana, Illinois.
- Haaland, D.M., Thomas, E.V. 1988. Partial Least-Squares Methods for Spectral Analyses. Relation to Other Quantitative Calibration Methods and the Extraction of Qualitative Information. *Analytical Chemistry*, 60: 1193-1202.
- Hammond, E.G., and Fehr, W.R. 1982. Progress in Breeding for Soybean Oil with Low Linolenic Acid. *J. Am. Oil Chem. Soc.* 59 (4): 304A.
- Hammond, E.G., and Fehr, W.R. 1983. Registration of A6 Germplasm Line of Soybean. *Crop Sci.* 23: 192-193.
- Harrick, N. J. 1960. Surface Chemistry from Spectral Analysis of Totally Internal Reflected Radiation. *J. Phys. Chem.* 64:1110.
- Haswell, S.J. 1992. *Practical Guide to Chemometrics*. Marcel Dekker, Inc., New York.
- Harris, Robin K. 1983. *Nuclear Magnetic Resonance spectroscopy, A Physicochemical View*, Pitman, London.
- Howard Mark. 2001. *Near-Infrared Applications in Biotechnology, (Biochemical and Pharmaceutical)*. Chapter 11 "Fundamentals of Near-Infrared Spectroscopy", P. 312, Edited by Raghavachari, R. Promega Corporation, Madison, Wisconsin.
- Hymowitz, T., Collins, F.I., Panczner, J. and Walker, W.M. 1972. Relationship between the Content of Oil, Protein, and Sugar in Soybean Seed. *Agron. J.* 64: 613-616.
- Isaksson, T., Miller, C.E. and Naes, T. 1992. Nondestructive NIR and NIT Determination of Protein, Fat and Water in Plastic Wrapped, Homogenized Meats. *Applied Spectroscopy*. 46: 1685.
- Isaksson, T., Naes, T. 1988. The Effect of Multiplicative Scatter Correction (MSC) and Linearity Improvement in NIR Spectroscopy. *Applied Spectroscopy*. 42(7): 1273-1284.
- Jones, D.B. 1931. Factors for Converting Percentages of Nitrogen in Foods and Feeds into Percentages of Nitrogen , USDA Circular No. 183, Washington DC.
- Joslyn, M.A. 1970. *Methods in Food Analysis*, 2nd ed. Academic Press, New York.
- Kabelka, E.A., Diers, B.W., Fehr, W.R., LeRoy, A.R., Baianu, I.C., You, T., Neece, D. J., Nelson, R. L. 2004. Putative Alleles for Increased Yield from Soybean Plant Introductions. *Crop Sci.*, 44:784-791.
- Kakalis, L.T. and Baianu, I.C. 1989. A High-Resolution ¹³C NMR Study of Glycinin in D₂O Solutions. *J. of Agric. & Food Chem.* 37: 1222-1228.
- Kakalis, F.T. and Baianu, I.C. 1990. High-Resolution ¹³C NMR Study of the Soybean 7S Storage Protein Fraction in Solution. *J. of Agric. & Food Chem.* 38: 1314-1320.
- Kettling, U., Koltermann, A., Schwille, P., and Eigen, M. 1998. Real-Time Enzyme Kinetics Monitored by Dual-Color Fluorescence Cross-Correlation Spectroscopy. *Proc. Natl. Acad. Sci. USA* 95: 1416-1420.
- Klempir J. M. 1999. Nuclear Magnetic Resonance Characterization of Soybeans for Oil Content Measurement. MS Thesis, University of Illinois at Urbana-Champaign, Urbana, Illinois.
- Kortum, G. 1969. *Reflectance Spectroscopy--Principles, Methods, Applications*. Springer-Verlag, NewYork, NY.

- Lamb, D., Hurburgh, C.R. 1991. Moisture Determination in Single Soybean Seeds by Near-Infrared Transmittance. *Trans. ASAE*. 34: 2123-2129.
- Lee, S.J., Ahn, J.K., Kim, S.H., Kim, J.T., Han, S.J., Jung, M.Y. and Chung, I.M. 2003. Variation in Isoflavone of Soybean Cultivars with Location and Storage Duration. *J. Agric. Food Chem.* 51(11): 3382-3389.
- Liu, K.S. 1997. *Soybeans: Chemistry, Technology, and Utilization*. Chapman & Hall, New York, NY. 532p.
- Liu, K.S. 2000. Expanding Soybean Food Utilization. *Food Technology*. 54(7): 46-58.
- Lowry, O. H., N.J. Rosebrough, A. L. Farr and R. J. Randall, 1951. Protein Measurement with the Folin Phenol Reagent. *J Biol. Chem.* 193: 265-275.
- Madden, H.M., 1978. Comments on the Savitzky-Golay Convolution Method for Least-Squares Fit Smoothing and Differentiation of Digital Data. *Anal. Chem.* 50: 1383.
- Maluszynski, M., Ahloowalia, B.S. and Sigurbjornsson, B. 1995. Application of *in vivo* and *in vitro* Mutation Techniques for Crop Improvement. *Euphytica*, 85: 303-315.
- Mansur, L.M., Orf, J.H., Chase, K., Jarvik, T., Cregan, P.B. and Lark, K.G. 1996. Genetic Mapping of Agronomic Traits Using Recombinant Inbred Lines of Soybean. *Crop Sci.* 36: 1327-1336.
- Mark, H. 2001. *Near-Infrared Applications in Biotechnology*, Chapter 11 "Fundamentals of Near Infrared Spectroscopy", p.312, Editor Raghavachari, R. Promega Corporation, Madison, Wisconsin.
- Martens, H., Jensen, S.A. and Geladi, P. 1983. Proceedings, Nordic Symposium on Applied Statistics, Stavanger, June 1983, pp.205-234.
- Mie, G. 1908. *Ann. Physik.* 4: 25, 377.
- Minson, D.J., Tuckett, P.G, Law D.P. 1985. A Comparison of three Methods of Increasing the Precision of Regression Used for Estimating Nutritive Value of Forages by Near Infrared Reflectance. 1985, NIR84 Proc. Int. Symp. Near Infrared Reflectance Spectroscopy., 1st. RACI, Cereal Chem. Div. Melbourne, Australia. pp 138-148.
- Myoung, G.C., In, Y.B., Sung, T.K., Won, Y.H., Doo, C.S., Huhn, P.M. and Kwang, H.K. 2001A. Determination of Protein and Oil Contents in Soybean Seed by Near Infrared Reflectance Spectroscopy. *Korean J Crop Sci.* 46(2): 106-111.
- Myoung, G.C., In, Y.B., Sung, T.K., Won, Y.H., Doo, C.S., Huhn, P.M. and Kwang, H.K. 2001B. Non-Destructive Method for Selection of Soybean Lines Contained High Protein and Oil by Near Infrared Reflectance Spectroscopy. *Korean J Crop Sci*, 46(5): 401-406.

- Nieh C.D. and Snyder, H.E. 1991. Solvent Extraction of Oil from Soybean Flour: I. Extraction Rate, a Countercurrent Extraction System, and Oil Quality. *J. Am. Oil Chem. Soc.* 68(4): 246-249.
- Nielsen, S.S. 1994. *Introduction to the Chemical Analysis of Foods*. Jones and Bartlet Publishers, Boston, London, Chapters 7, 10, 14.
- Norris, K.H. and Williams, P.C. 1984. Optimization of Mathematical Treatment of Raw Near-IR Signal in the Measurement of Protein in Hard Red Spring Wheat: 1. Influence of Particle Size. *Cereal Chem.* 62: 158-165.
- Norris, K.H. 1983. Using Gap Derivatives as Pre-processing for Quantitative Models, Food Research & Data Analysis, Proceedings of the 1982 IUFST Symposium, Martens, H. Ed., Applied Science Publishers, Oslo, 46-47.
- Ohnishi, S.T. and Barr, J.K. 1978. A Simplified Method of Quantitating Protein Using the Biuret and Phenol Reagents. *Anal. Biochem.* 86: 193-200.
- Openshaw, S.J. and Hadley, H.H. 1984. Selection Indexes to Modify Protein Concentration of Soybean Seeds. *Crop Sci.* 24: 1-4.
- Orman, B.A., Schumann, R.A., 1992. Nondestructive Single-Kernel Oil Determination of Maize by Near-Infrared Transmission Spectroscopy. *J. Am. Oil Chem. Soc.* 69: 1036-1038.
- Osborne, B. G., Fearn, T. and Hindle, P.H. 1993. *Practical NIR Spectroscopy with Applications in Food and Beverage Analysis*, 2nd Edition. pp 13-21, 99-117. Longman Scientific & Technical, Harlow, UK. 227p.
- Paulsen, M.R., Bajaj, M. 2000. Using Unscrambler for Extractable Corn Starch Calibration. ASAE Paper No. 00-3091. ASAE. St. Joseph, MI.
- PerkinElmer Instruments, 2000. Spectrum One User's Guide, Shelton, CT 06484, USA.
- PerkinElmer Instruments, 2002. AutoImage User's Guide, Sheldon, CT 06484, USA.
- Peterson, G. L. 1979. Review of the Folin Phenol Protein Quantitation Method of Lowry, Rosebrough, Farr and Randall. *Analytical Biochemistry* 100: 201-220.
- Pfeffer, P. and Gerasimowicz, W. (Eds.). 1987. *NMR in Agriculture*. CRS Publishers, Boca Raton, FL.
- Pomeranz Y. and Meloan, C.E. 1994. *Food Analysis Theory and Practice*, 3rd edition, chapter 33, 567-574. Chapman & Hall, New York, NY.

- Raghavachari, R. 2001. *Near-Infrared Applications in Biotechnology*, Marcel-Dekker, New York, NY.
- Reffer, J.A., Martoglio, P.A. 1995. *Uniting Microscopy and Spectroscopy, Practical Guide to Infrared Microspectroscopy*, Chapter 2, p.41, Edited by Howard J. Humecki, Marcel Dekker, New York, NY.
- Rigler R. and Widengren J. 1990. Ultrasensitive Detection of Single Molecules by Fluorescence Correlation Spectroscopy, *BioScience* (Ed. Klinge & Owman) p.180.
- Rippe, K. 2000. Simultaneous Binding of Two DNA Duplexes to the NtrC-Enhancer Complex Studied by Two-Color Fluorescence Cross-Correlation Spectroscopy. *Biochemistry* 39(9): 2131-2139.
- Robertson, J. A. and F. E. Barton.1984. Oil and Water Analysis of Sunflower Seeds by Near Infrared Reflectance spectroscopy, *JAOCS*, 61 (3): 543-547.
- Robinson, H.W. and C.G. Hodgen 1940. The Biuret Reaction in the Determination of Serum Protein. 1. A Study of the Conditions Necessary for the Production of the Stable Color which Bears a Quantitative Relationship to the Protein Concentration. *Journal of Biological Chemistry* 135: 707-725.
- Rutar, V. 1982. A New Possibility for Nondestructive Protein Content Determination in Viable Seeds, *Applied Spectroscopy*, 36: 259-260.
- Rutar, V., M. Kovac and G. Lahajnar.1989. Nondestructive Study of Liquids in Single Seeds Using Nuclear Magnetic Resonance and Magic Angle Sample Spinning, *JAOCS*, 66: 961-965.
- Rutar, V. 1984. NMR Studies in Intact Seeds, in *Nuclear Magnetic Resonance in Agriculture*. editors Philip E. Pfeffer and Walter V. Gerasimowicz, CRC Press Inc., Boca Raton, Florida.
- Rutar, V.1989. Magic Angle Spinning NMR Spectroscopy of Liquids as Nondestructive Method for Studies of Plant Seeds, *J. Agric. Food Chem.*, 37: 67-70.

- Savitsky, A. and Golay, M.J. 1964. Smoothing and Differentiation of Data by Simplified Least Squares Procedures. Anal. Chem. 36: 1627-1639.
- Schwille P., Oehlenschläger F. and Walter N. 1996. Analysis of RNA-DNA Hybridization Kinetics by Fluorescence Correlation Spectroscopy, Biochemistry 35: 10182.
- Schwille, P., Bieschke, J. and Oehlenschläger F. 1997. Kinetic Investigations by Fluorescence Correlation Spectroscopy: The Analytical and Diagnostic Potential of Diffusion Studies, Biophys. Chem. 66: 211-228.
- Schwille P., Meyer-Almes F.-J., and Rigler R. 1997. Dual-Color Fluorescence Cross-Correlation Spectroscopy for Multicomponent Diffusional Analysis in Solution, Biophys. J. 72: 1878-1886.
- Schwille, P., Haupts, U., Maiti, S., and Webb. W. 1999. Molecular Dynamics in Living Cells Observed by Fluorescence Correlation Spectroscopy with One- and Two-Photon Excitation. Biophysical Journal, 77(10):2251-2265.
- Schwille, P. 2001. *Fluorescence Correlation Spectroscopy. Theory and applications*. Rigler R., and Elson, E. S. eds. Springer Verlag: Berlin. p360.
- Shchegolikhin, A., Lazareva, O. 2000. Diffuse Reflectance for the Routine Analysis of Liquids and Solids. Internet Journal of Vibration Spectroscopy [www.ijvs.com]. 4(1): 8-38.
- Shadow, W. 1998. *Rapid Analysis for the Food Industry Using Near-Infrared Spectroscopy*. Perten Instruments North America, Inc. 17p.
- Shadow, W. and A. Carrasco. 2000. Practical Single-Kernel NIR/Visible Analysis for Small Grains. Cereal Foods World 45(1): 16-18.
- Silvela, L., Rodgers, R., Barrera, A., Alexander, D.E. 1989. Effect of Selection Intensity and Population Size on Percent Oil in Maize, Zea mays L. Theoretical and Applied Genetics 78: 298-304.
- Simpson, A.M. Jr. and Wilcox, J.R. 1983. Genetic and Phenotypic Associations of Agronomic Characteristics in Four High Protein Soybean Populations [Glycine max]. Crop Sci. 23: 1077-1081.

- Smith, A.L. 1979. Ed. *Applied Infrared Spectroscopy: Fundamentals, Techniques, and Analytical Problem-solving*. John Wiley & Sons. New York, NY.
- Smith, P.K., Krohn, R.I., Hermanson, G.T., Mallia, A.K., Gartner, F.H., Provenzano, M.D., Fujimoto, E.K., Goeke, N.M., Olson, B.J., and Klenk, D.C. 1985. Measurement of Protein Using Bicinchoninic Acid. *Anal. Biochem.* 150: 76-85.
- Song, T., Barua, K., Buseman, G. and Murphy, P.A. 1998. Soy Isoflavone Analysis: Quality Control and a New Internal Standard. *Am. J. Clinical Nutrition.* 68(6), Supplement, 1474S-1479S.
- Sorvaniemi, J., Kinnunen, A., Tsados, A., Malkki, Y. 1993. Using Partial Least Squares Regression and Multiplicative Scatter Correction for FT-NIR Data Evaluation of Wheat Flours, *Lebensmittel-Wissenschaft & Technologie*, 26(3): 251-258.
- Specht, J., Nelson, R.L., Webster, R., Naidu, S., Ainsworth, E. and Ort, D. 2002. A Genomic Perspective on the Soybean Protein(+)/Oil(-)/Yield(-) Enigma. Program and Proceedings for the 9th Biennial Conference of the Cellular and Molecular Biology of the Soybean, August 11-14, 2002, P.202.
- Stuber, C.W. 1992, Biochemical and Molecular Markers in Plant Breeding. *Plant Breeding Rev.* 9: 37-61.
- Tajuddin, T., Watanabe, S., Masuda, R., Harada, K. and Kawano, S. 2002. Application of Near Infrared Transmittance Spectroscopy to the Estimation of Protein and Lipid Contents in Single Seeds of Soybean Recombinant Inbred Lines for Quantitative Trait Loci Analysis. *J. Near Infrared Spectrosc.* 10(4): 315-325.
- Tekel, J., Daeseleire, E., Heeremans, A. and Peteghem, C.V. 1999. Development of a Simple Method for the Determination of Genistein, Daidzein, Biochanin A, and Formononetin (Biochanin B) in Human Urine. *J. Agric. Food Chem.* 47(9): 3489-3494.
- Theissing, H. H. 1950. *J. Opt. Soc. Am.* 40: 232.
- Thompson, J.A. and Nelson, R.L. 1998. Utilization of Diverse Germplasm for Improving Soybean Yield. *Crop Sci.* 38: 1362-1368.
- Thompson N.L. 1991. In *Topics of Fluorescence Spectroscopy*. Lakowicz, J.R. ed. Plenum Press, Vol 1 p 337.
- Tiwari, P. N. and Gambhir, P. N. 1995. Seed Oil Determination without Weighing and Drying the Seeds by Combined Free Induction Decay and Spin-Echo Nuclear Magnetic Resonance Signals, *JAOCS*, 72(9).
- USDA National Agriculture Statistical Service (NASS). 1999. USDA-NASS Agricultural Statistics 1999. <http://www.usda.gov/nass/pubs/agr99/acro99.htm>.
- Velasco, L., Goffman, F.D., and Becker H.C. 1999. Development of Calibration Equations to Predict Oil Content and Fatty Acid Composition in Brassicaceae Germplasm by Near Infrared Reflectance Spectroscopy. *JAOCS*, 76: 25-30.
- Verma, D.P.S., Shoemaker, R.C. 1996. *Soybean Genetics, Molecular Biology and Biotechnology*. Cab International. Guildford, UK.

- Watson et al. 1987. Ed. *Corn: Chemistry and Technology*. American Association of Cereal Chemists. Inc. St. Paul, MN.
- Wang, D., Dowell, F.E., Lacey, R.E. 1999. Single Wheat Kernel Color Classification Using Near-Infrared Reflectance Spectra. *Cereal Chem.* 76: 30-33.
- Wehrmann, V.K., Fehr, W.R., Cianzio, S.R. and Cavins, J.F. 1987. Transfer of High Seed Protein to High Yielding Soybean Cultivars. *Crop Sci.* 25: 927-931.
- Wesley, I.J., Osborne, B.G., Anderssen, R.S and Skerritt, J.H. 1999. Curve Fitting Applied to Near Infrared Deconvolution of Wheat Functional Proteins in Flour. *Near Infrared Spectroscopy, Proceedings of the International Conference, 9th, Verona, Italy*, p.215.
- Whistler, R. L. and BeMiller, J. N. 1997, *Carbohydrates Chemistry for Food Scientists*, Eagen Press, St. Paul, MN.
- Wilcox, J.R., Cavins, J.F., and Nielsen, N.C. 1984. Genetic Alteration of Soybean Oil Composition by a Chemical Mutagen. *Journal of the American Oil Chemists Society*, 61: 97-100.
- Wilcox, J.R., Cavins, J.F. 1995. Backcrossing High Seed Protein to a Soybean Cultivar. *Crop Science*. 35: 1036-1041.
- Wilcox, J.R. 1998. Increasing Seed Protein in Soybean with Eight Cycles of Recurrent Selection. *Crop Sci.* 38, 1536-1540.
- Wilcox J.R. and Shibles, R.M. 2001. Interrelationship among Seed Quality Attributes in Soybean, *Crop Sci.* 41: 11-14.
- Williams, P.C. 1975. Application of NIRS to Analysis of Cereal and Oilseeds. *Cereal Chem.* 52:561-576.
- Williams, P.C., Sobering, D.C. 1993. Comparison of Commercial Near Infrared Transmittance and Reflectance Instruments for Analysis of Whole Grains and Seeds. *J. Near Infrared Spectroscopy*. 1: 25-32.

- Williams, P.C., Norris, K. 1987. *Near-Infrared technology in the agricultural and food industries*. American Association of Cereal Chemists, Inc. St.Paul, MN. p330.
- Wilson, J. M., Kramer, A., and Ben-Gera, I. 1973. Quantitative Determination of Fat, Protein and Carbohydrates of Soy Products with Infrared Attenuated Total Reflectance. *J. Food Sci.* 38:14-17.
- Wilson, R. F. Advances in the Genetic Alteration of Soybean Oil Composition. 1991. In: R.F. Wilson (ed). *"Designing Value-Added Soybean for Markets of the Future."* American Oil Chemists Society, Champaign, Illinois p.38-53.
- Winkler, T., Bieschke, J. and Schwill, P. 1997. Development of a Dual-Color Cross-Correlation System for FCS. www.mpibpc.gwdg.de/abteilungen/081/fcs/correlation/english.html.
- Winkler, T., Kettling, U., Koltermann, A., and Eigen, M. 1999. Confocal Fluorescence Coincidence Analysis: An Approach to Ultra High-Throughput Screening. *Proc. Natl. Acad. Sci. USA*, 96: 1375-1378.
- You, T. PhD Thesis, 2005, University of Illinois, Urbana, Illinois.
- You, T., Guo J., Baianu, I.C., 1999. Soybean Composition Analysis by Near Infrared Spectroscopy. *Global Soy1999*, Chicago, Illinois(Abtract).
- You, T., Guo, J., Baianu, I.C. and Nelson, R.L. 2002A. Determination of Isoflavones Contents for Selected Soybean Lines by Fourier Transform Near Infrared Reflectance Spectroscopy. *Proceedings for the 9th Biennial Conference of the Cellular and Molecular Biology of the Soybean*, August 11-14, 2002, p505.
- You T., Guo, J., Baianu, I.C., Nelson, R.L. 2002B. Rapid Determination of Protein, Oil, Moisture, and Isoflavones Contents of Single Soybean Seeds by Fourier Transform Near Infrared Reflectance Spectroscopy. *Proceedings of the China and International Soy Conference and Exhibition 2002 (CISCE 2002)*, November 6-9, 2002, 414-415.

8. SPECIALIZED REFERENCES

1. Silvela, L., Rodgers, R., Barrera, A., Alexander, D.E. (1989) *Ther. Appl. Genet.* 78:298
2. Lowry, O.H., Rosebrough, N.J., Farr, A.L., and Randall, R.J. (1951) Protein measurement with the Folin phenol reagent. *J. Biol Chem.* 193, 265-275.
3. Williams, P.C. (1975) *Cereal Chem.* 52, 561-576.
4. Ben-Gera, I., Norris, K.H. (1968) *Isr. J. Agric. Res.* 18, 125-132.
5. Shadow, W. (1998) Rapid analysis for the food industry using near-infrared spectroscopy. Perten Instruments North America, Inc.
6. Lamb, D., Hurburgh, C.R. (1991) *Trans. ASAE.* 34, 2123-2129.
7. Orman, B.A., Schumann, R.A. (1992) *JAOCs.* 69, 1036-1038.
8. Wang, D., Dowell, F.E., Lacey, R.E. (1999) *Cereal Chem.* 76, 30-33.
9. Shadow, W. and Carrasco, A. (2000) Practical single-kernel NIR/visible analysis for small grains. *Cereal Foods World.* 45, 16-18.
10. Raghavachari, R. (2001) *Near-Infrared Applications in Biotechnology*, Marcel-Decker, New York, NY p 125.
11. Barton, F.E. (2002) *Spectroscopy Europe.* 14, 12-18.
12. Baianu, I.C., and Kumosinski, T.F. (1993) In *Physical Chemistry of Food Processes. Vol.2. Advanced Techniques, Structures, and Applications.* Van Nostrand Reinhold. New York, NY.
13. Bloch, F. (1946) Nuclear Induction. *Phys. Rev.* 70, 460-474.
14. Abragam, A. (1961) *The Principles of Nuclear Magnetism*, Clarendon Press, Oxford.
15. Slichter, C.P. (1963) *Principles of Magnetic Resonance.* Harper and Row Press. New York.
16. AOAC International. (1995). *Official Methods of Analysis*, 16th ed., AOAC International, Gaithersburg, MD.
17. Becker, E.D. (1980) *High Resolution NMR. Theory and Chemical Applications*, 2nd Ed. New York : Academic Press
18. Mansfield, P. (1965) *Physics Review.* A137, 961.

19. Baianu, I.C., You, T., Guo, J., Nelson, R.L. (2002) Calibration of Dual Diode Array and Fourier Transform Near Infrared Reflectance Spectrometers for Composition Analysis of Single Soybean Seeds in Genetic Selection, Cross-Breeding Experiments. *Soybean2002 Conference*, Urbana, IL.
20. Williams, P., and Norris, K. (1987) *Near-Infrared technology in the agricultural and food industries*. American Association of Cereal Chemists, Inc. St.Paul, MN.
21. You, T., Guo, J., Baianu, I.C., Nelson, R.L. (2003) Diode-Array Near Infrared Spectroscopy for Rapid Soybean Composition Analysis: Light Scattering Corrections for Intact and Ground Soybean Seeds. (*In submission*)
22. Feynman, R.P. (1963) *Lectures on Physics Vol 3*. Addison-Wesley Publishing Company, Reading, MA.
23. Kortum, G. (1969) *Reflectance Spectroscopy-Principles, Methods, Applications*. Springer-Verlag, NewYork.
24. Mie, G. (1908) *Ann. Physik*. 25, 377.
25. Theissing, H. H. (1950) *J. Opt. Soc. Am.* 40, 232.
26. You, T., Guo, J., Baianu, I.C., and Nelson, R.L (2002). Rapid Determination of Protein, Oil, Moisture, and Isoflavones Contents of Single Soybean Seeds by Fourier Transform Near Infrared Reflectance Spectroscopy. *China & International Soybean Conference & Exhibition 2002*, Beijing, China.
27. Geladi, P., MacDougall, D., Martens, H. (1985) Linearization and Scatter-Correction for Near-Infrared Reflectance Spectra of Meat. *Applied Spectroscopy*. 39(3), 491-500.
28. Isaksson, T., and Naes, T. (1988) The effect of Multiplicative Scatter Correction (MSC) and Linearity Improvement in NIR Spectroscopy. *Applied Spectroscopy*. 42(7), 1273-1284.
29. Haaland, D.M., and Thomas, E.V. (1988) *Analytical Chemistry*. 60, 1193.
30. Sorvaniemi, J., Kinnunen, A., Tsados, A., and Malkki, Y. (1993) Using partial least squares regression and multiplicative scatter correction for FT-NIR data evaluation of wheat flours. *Lebensmittel-Wissenschaft & Technologie* 26(3), 251-258.
31. AOCS. (2002) Additions and Revisions to the Official Methods and Recommended Practices of the AOCS. *AOCS Press*. Champaign, IL. AK4-95.
32. Baianu, I.C. (1989) In *NMR in Agriculture*. Pfeffer, P. and Gerasimowicz. W., eds., CRC Press. Boca Raton, FL.
33. Ohnishi, S.T., and Barr, J.K. (1978) A simplified method of quantitating protein using the biuret and phenol reagents. *Anal. Biochem*. 86, 193-200.

34. Anderson, R., Killan, A.S., and Orf, J.H. (2002) Amino Acid Analysis of Soybean Powders by NIR. Soy 2002 Conference (Abstr.)
35. Bardet, M., Foray, M.F., Bourguignon, J., and Krajewski P. (2001) Investigation of seeds with high-resolution solid-state ^{13}C NMR. *Magnetic Resonance in Chemistry*. 39, 733-738.
36. Kakalis, L.T., and Baianu, I.C. (1989) A high-resolution ^{13}C NMR study of glycinin in D_2O solutions. *J. of Agric. & Food Chem.* 37, 1222-1228.
37. Kakalis, F.T., and Baianu, I.C. (1990) High-resolution ^{13}C NMR study of the soybean 7S storage protein fraction in solution. *J. of Agric. & Food Chem.* 38, 1314-1320.
38. Baianu, I.C. (1981) Carbon-13 and Proton NMR studies of Wheat Proteins in Solution. *J. Sci. of Food & Agriculture*. 32, 309-313.
39. Baianu, I.C. (1982) High Resolution Carbon-13 and Nitrogen-15 NMR Studies of Flinor Wheat Gliadins: Spectral Assignments, Changes with concentration and heat treatments. *J. Sci. of Food & Agriculture*. 33, 373-383.
40. Augustine, M., and Baianu, I.C. (1986) High Resolution Carbon-13 NMR Studies of Maize Proteins. *J. Cereal Sci.* 4, 371-378.
41. Baianu, I.C. (1987) 125 MHz Carbon-13 NMR Spectra of Corn Zeins in Alkaline Solutions. Unpublished results.
42. Baianu, I.C. (1990) In NMR Applications in Biopolymers. Finley J. W. Ed. Plenum Press, New York.
43. Baianu, I.C., and Förster H. (1980) Cross-polarization, high-field Carbon-13 NMR techniques for studying Physicochemical Properties of Wheat Grains and Wheat Proteins., *J. Applied Biochemistry*. 2, 347-355.
44. Baianu, I.C., et al.(1990) In *NMR Applications in Biopolymers..* Finley, J.W. ed. Plenum Press, New York.
45. Wei, T.C. (1990) Molecular Dynamics, Conformation and Hydration Studies of Soybean Proteins and Amino Acids by NMR. Ph.D. Thesis, University of Illinois, Urbana.
46. Augustine, M., and Baianu, I.C. (1984) High-resolution ^{13}C NMR and Ion Exchange Chromatography Amino Acid Profiling of Corn and Soybean Protein Hydrolizates. *FASEB Proceed.* 1125.
47. Rutar, V. (1989). *J. of Agric. And Food Chem.* 37, 67.
48. Levitt, M.H. (2001) *Spin Dynamics: Basics of Nuclear Magnetic Resonance*. John Willey & Sons, New York.

49. Baianu, I.C. (1992) *Physical Chemistry of Food Processes*. Vol. 1: *Fundamental Aspects*. Van Nostrand Reinhold. New York, NY.
50. Velasco, L., Goffman, F.D., and Becker H.C. (1999). Development of Calibration Equations to Predict Oil Content and Fatty Acid Composition in *Brassicaceae* Germplasm by Near- Infrared Reflectance Spectroscopy. *JAOCs*, 76, 25-30.
51. Sato, T., (2002). New Estimation Method for Fatty Acid Composition in Oil using Near Infrared Spectroscopy. J generation systems. *Biosci. Biotechnol. Biochem.* 66, 2543-2548.
52. Pazdernik, D.L., Killam, A. S., and Orf, J. H. (1997) *Agronomy Journal*. 89, 679-685.
53. Windham, W.R., Mertens, D.R., and Barton F.E. (1989) Protocol for NIRS calibration: Sample selection and equation development and validation. In Near Infrared reflectance spectroscopy (NIRS): *Analysis of forage quality*. *Agric. Handb. 643*. USDA-ARS, Washington, DC.
54. Guo, J., You, T., Baianu, I.C. and Nelson, R. (2002) Development of calibrations for Bulk Whole and Ground Yellow Soybean Seeds obtained by Diode Array Near Infrared Reflectance Spectroscopy. *International Soy Conference and Exhibition*. Beijing, China.
55. Thompson, J.A. and Nelson, R.L. (1998) Utilization of diverse germplasm for improving soybean yield. *Crop Sci.* 38, 1362-1368.
56. Kabelka, E. A., Diers, B. W., Fehr, W. R. , LeRoy, A.R., Baianu, I.C., You, T., Neece, D.J., and Nelson, R.L. (2003) Identification of putative yield enhancing quantitative trait loci from exotic soybean germplasm. *Crop Sci.* (In Press)
57. Wilcox, J. R. and Cavins, J. F. (1995) Backcrossing high seed protein to a soybean cultivar. *Crop-Sci*, 35(4), 1036-1041.

FCS and FCCS REFERENCES

1. Eigen, M., and Rigler, R. (1994) Sorting single molecules: Applications to diagnostics and evolutionary biotechnology, *Proc. Natl. Acad. Sci. USA* **91**, 5740.
2. Winkler, T., Bieschke, J. and Schwille, P. (1997) Development of a dual-color cross-correlation system for FCS.
www.mpibpc.gwdg.de/abteilungen/081/fcs/correlation/english.html.
3. Schwille, P., Haupts, U., Maiti, S., and Webb, W. (1999). Molecular dynamics in living cells observed by fluorescence correlation spectroscopy with one- and two-photon excitation. *Biophysical Journal* **77**(10), 2251-2265.
4. Chen, Y., Müller, J.D, Tetin, S.Y., Tyner, J.D., and Gratton, E. (2000) *Biophys. J.* **79**, 1074
5. Bagatolli, L.A., and Gratton, E. (2000) A correlation between lipid domain shape and binary phospholipid mixture composition in free standing bilayers: a two-photon fluorescence microscopy study. *Biophys J.* **79**, 434-447.
6. Bagatolli, L.A., and Gratton, E. (2000) Two-photon fluorescence microscopy of coexisting lipid domains in giant unilamellar vesicles of binary phospholipid mixtures. *Biophys J.* **78**, 290-305.
7. Murashige, T. and Skoog, F. (1962) A revised medium for rapid growth and bioassay with tobacco tissue cultures. *Physiol. Plant.* **15**, 473-497.
8. Gamborg, O.L., Miller, R.A., and Ojima, K. (1968) Nutrient requirement of suspension cultures of soybean root cells. *Exp. Cell Res.* **50**, 151-158.
9. Hofmann, N.E, Raja, R., Nelson, R.L., and Korban, S.S. (2003) Mutagenesis of embryogenic cultures of soybean and detecting polymorphisms using RAPD markers. *Biol. Plant.* (in press).

10. Clarke, F.C., Jee, D.R., Moffat, A.C., and Hammond, S.V. (2001) Effective sample volume for measurements by Near-Infrared Microscopy. (e.g.: 40,000 m³, or ~30 ng); (Abstract, 116), *British Pharmaceutical Conference*. UK.
11. Baianu, I.C. (1996) NMR Principles and Applications to the Structure and Hydration of Food Systems, *Physical Chemistry of Food Processes*. Vol 2. Van Nostrand Reinhold, New York, p 338.
12. Diaspro, A., and Robello, M. (1999) Multi-photon Excitation Microscopy to Study Biosystems. *European Microscopy and Analysis*, 5, 5-7.
13. Rigler R., Mets Ü., Widengren J. and Kask P. (1993). Fluorescence correlation spectroscopy with high count rate and low background: Analysis of translational diffusion, *Eur. Biophys. J.* 22,169.
14. Chirico, G., Bettati, S., Mozzarelli, A., Chen, Y., Müller, J. D., and Gratton, E. (2001). Molecular Heterogeneity of O-Acetylserine Sulfhydrylase by Two-Photon Excited Fluorescence Fluctuation Spectroscopy. *Biophys J*, 80(4), 1973-1985.
15. Baianu, I.C., Costescu, D.M., and You, T. (2002). Novel Techniques for Microspectroscopy and Chemical Imaging Analysis of Soybean Seeds and Embryos. *Soy2002 Conference*, Urbana, Illinois.
16. Baianu, I.C., Costescu, D.M., Hofmann, N., and Korban, S.S. (2003). Near Infrared Microspectroscopy, Chemical Imaging and NMR Analysis of Oil in Developing and Mutagenized Soybean Embryos in Culture. *Poster, AOCs Meeting, Analytical Division*.
17. You, T., Guo, J., Baianu, I.C. and Nelson, R.L. (2002). Determination of Isoflavone Contents for Selected Soybean Lines by FT-NIR Reflectance Spectroscopy. *Soy2002 Conference*. Urbana, Illinois.
18. Thompson N.L. (1991) In Topics of Fluorescence Spectroscopy. Lakowicz, J.R. ed. Plenum Press, Vol 1 p 337.

19. Schwille, P., Bieschke, J. and Oehlenschläger F. (1997) Kinetic investigations by fluorescence correlation spectroscopy: The analytical and diagnostic potential of diffusion studies, *Biophys. Chem.* 66, 211-228.
20. Schwille P., Oehlenschläger F. and Walter N. (1996) Analysis of RNA-DNA hybridization kinetics by fluorescence correlation spectroscopy, *Biochemistry* 35, 10182.
21. Walter N., Schwille P. and Eigen M. (1996). Fluorescence correlation analysis of probe diffusion simplifies quantitative pathogen detection by PCR, *Proc. Natl. Acad. Sci. USA* 93,12805.
22. Oehlenschläger F., Schwille P. and Eigen M. (1996). Detection of HIV-1 RNA by nucleic acid sequence-based amplification combined with fluorescence correlation spectroscopy, *Proc. Natl. Acad. Sci. USA* 93, 1281.
23. Winkler, T., Schwille, P., and Oehlenschläger, F. (1998). Detection of HIV-1 RNA by NASBA-FCS. <http://www.mpibpc.gwdg.de/abteilungen/081/fcs/nasba/english.html>.
24. Compton, J. (1991). Nucleic acid sequence-based amplification. *Nature* 350, 91-92.
25. Schwille, P., Oehlenschläger, F. and Walter, N.G. (1997). Comparative hybridization kinetics of DNA-oligonucleotides to a folded RNA target in solution. A summary of measurements and results based on FCS diffusion analysis was presented by: Schwille P., Bieschke J. and Oehlenschläger F. In: Kinetic investigations by fluorescence correlation spectroscopy: The analytical and diagnostic potential of diffusion studies. *Biophys. Chem.* 66, 211-228.
26. Kettling, U., Koltermann, A., Schwille, P., and Eigen, M. (1998) Real-time enzyme kinetics monitored by dual-color fluorescence cross-correlation spectroscopy. *Proc. Natl. Acad. Sci. USA* 95, 1416-1420.
27. Koltermann, A., Kettling, U., Bieschke, J., Winkler, T., and Eigen, M. (1998) Rapid assay processing by integration of dual-color fluorescence cross-correlation spectroscopy: High throughput screening for enzyme activity *Proc. Natl. Acad. Sci. USA* 95, : 1421-1426.

28. Schwille P., Meyer-Almes F.-J., and Rigler R. (1997). Dual-color fluorescence cross-correlation spectroscopy for multicomponent diffusional analysis in solution, *Biophys. J.* 72, 1878.
29. Winkler, T., Kettling, U., Koltermann, A., and Eigen, M. (1999) Confocal fluorescence coincidence analysis: An approach to ultra high-throughput screening *Proc. Natl. Acad. Sci. USA* 96,: 1375-1378.
30. Rigler R. and Widengren J. (1990). Ultrasensitive detection of single molecules by fluorescence correlation spectroscopy, *BioScience (Ed. Klinge & Owman)* p.180.
31. Rippe, K. (2000) Simultaneous Binding of Two DNA Duplexes to the NtrC-Enhancer Complex Studied by Two-Color Fluorescence Cross-Correlation Spectroscopy. *Biochemistry* 39(9), 2131-2139.
32. Schwille, P. (2001) Fluorescence Correlation Spectroscopy. Theory and applications R. Rigler R., and Elson, E. S. eds. Springer Verlag: Berlin. p360.
33. Glover, P. and P. Mansfield. (2002). Limits to Magnetic Resonance Microscopy. *J. Rep. Prog. Phys.* 65, 1489-1511.
34. Lee, S. C. et al., (2001). One Micrometer Resolution NMR Microscopy. *J. Magn. Res.*, 150, 207-213.
35. Baianu, I.C. (1985). High-Field NMR of Food Systems and Grains. *ACS Symp.*, Washington D.C.
36. Otting, G., and Wutrich K. (1989) Studies of Protein Hydration in Aqueous Solution by Direct NMR Observation of Individual Protein-Bound Water Molecules. *J. Am. Chem. Soc.* 102, 3636-3637.

RECENT PREPRINTS:

<http://precedings.nature.com/documents/6200/version/1/files/npre20116200-1.pdf>

**Near Infrared Microspectroscopy, Fluorescence
Microspectroscopy, Infrared Chemical Imaging and
High-Resolution Nuclear Magnetic Resonance Analysis of Soybean
Seeds, Somatic Embryos and Single Cells**

**I.C. Baianu^{*1-3}, D. Costescu²⁻³, T. You¹⁻², P.R. Lozano¹⁻², N.E. Hofmann² and
S.S. Korban⁴**

<http://aux.planetphysics.us/files/lec/260/Soy2002-Imaging-2.pdf>

**Novel Techniques for
Microspectroscopy and Chemical
Imaging Analysis of Soybean
Seeds and Embryos**

I.C. Baianu^{*,†}, Tiefeng You[†], Doina Costescu⁺